

**Patient Information**  
**Natesto (Na-tes-to) CIII**  
**(testosterone) nasal gel**

**What is Natesto?**

- Natesto is a prescription medicine that contains testosterone and is used to treat adult males who have low or no testosterone due to certain medical conditions.
- Your healthcare provider will test the testosterone level in your blood before you start and while you are using Natesto.
- It is not known if Natesto is safe or effective to treat men who have low testosterone due to aging.
- It is not known if Natesto is safe or effective in children less than 18 years old. Improper use of Natesto may affect bone growth in children.
- Natesto is a controlled substance (CIII) because it contains testosterone that can be a target for people who abuse prescription medicines. Keep your Natesto in a safe place to protect it. Never give your Natesto to anyone else, even if they have the same symptoms you have. Selling or giving away this medicine may harm others and is against the law.
- Natesto is not meant for use in women.

**Who should not use Natesto?**

**Do not use Natesto if you:**

- are a man who has breast cancer
- have or might have prostate cancer
- are pregnant or may become pregnant or are breastfeeding. Natesto may harm your unborn or breastfeeding baby.

Talk to your healthcare provider before using Natesto if you have any of the above conditions.

**Before using Natesto, tell your healthcare provider about all of your medical conditions, including if you:**

- have or have had nose (nasal) or sinus problems **or** nasal or sinus surgery
- have had a broken nose (fracture) within the past 6 months
- have or have had a fracture of your nose that caused the inside of your nose to be crooked (deviated anterior nasal septum)
- have or have had problems with swelling of the lining of your nose (mucosal inflammatory disorder)
- have breast cancer
- have or might have prostate cancer
- have urinary problems due to an enlarged prostate
- have heart problems
- have liver or kidney problems
- have problems breathing while you sleep (sleep apnea)

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, or herbal supplements.

Using Natesto with certain other medicines can affect each other. Especially tell your healthcare provider if you take:

- insulin
- medicines that decrease blood clotting
- corticosteroids

**How should I use Natesto?**

- See the detailed Instructions for Use at the end of this Patient Information for Instructions about the right way to use Natesto.
- Use Natesto exactly as your healthcare provider tells you to. Your healthcare provider will tell you how much Natesto to use and when to use it.

- Natesto is for use in your nose (intranasally) only.
- Natesto can be used with a type of nasal spray called sympathomimetic decongestants such as oxymetazoline. Natesto should not be used with other nasal sprays. It is not known if Natesto is safe and effective when used with other nasal sprays.

**What are the possible side effects of Natesto?**

**Natesto may cause serious side effects, including:**

- **Nose (nasal) problems.** Signs and symptoms of nasal problems may include runny nose, congestion, sneezing, nose bleed, nasal discomfort, nasal scabbing, or nasal dryness.
- **If you already have enlargement of your prostate gland, your signs and symptoms can get worse while using Natesto.** This can include: Increased urination at night, trouble starting your urine stream, having to pass urine many times during the day, having an urge that you have to go to the bathroom right away, having a urine accident, being unable to pass urine or weak urine flow.
- **Possible increased risk of prostate cancer.** Your healthcare provider should check you for prostate cancer or any other prostate problems before you start and while you use Natesto.
- **Changes in red blood cells.**
- **Blood clots in your legs or lungs.** Signs and symptoms of a blood clot in your legs can include leg pain, swelling or redness. Signs and symptoms of a blood clot in your lungs can include difficulty breathing or chest pain.
- **Possible increased risk of heart attack or stroke.**
- **In large doses, Natesto may lower your sperm count.**
- **Possible increased risk of liver problems.** Signs and symptoms of liver problems may include: Nausea or vomiting, yellowing of your skin or whites of your eyes, dark urine, or pain on the right side of your stomach area (abdominal pain).
- **Swelling of your ankles, feet, or body, with or without heart failure.** This may cause serious problems for people who have heart, kidney, or liver disease.
- **Enlarged or painful breasts.**
- **Problems breathing while you sleep (sleep apnea).**

Call your healthcare provider right away if you have any of the serious side effects listed above.

These are not all the possible side effects of Natesto. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**The most common side effects of Natesto include:**

- increased prostate specific antigen (a test used to screen for prostate cancer)
- headache
- runny nose
- nose bleeds
- nose pain
- sore throat
- cough
- upper respiratory infection
- sinus infection
- nose scabs

**Other side effects include** more erections than normal for you or erections that last a long time.

**How should I store Natesto?**

- Store Natesto between 68°F to 77°F (20°C to 25°C).
- **Keep Natesto and all medicines out of the reach of children.**

**General information about the safe and effective use of Natesto.**

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use Natesto for a condition for which it was not prescribed. Do not give Natesto to other people, even if they have the same symptoms that you have. It may harm them.

If you would like more information about Natesto, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information about Natesto that is written for health professionals. For more information go to [www.natesto.com](http://www.natesto.com) or call 1-833-698-3786.

**What are the ingredients in Natesto?**

**Active ingredient:** testosterone

**Inactive ingredients:** castor oil, oleoyl polyoxylglycerides, and colloidal silicon dioxide

Marketed by: Acerus Pharmaceuticals Corporation., Mississauga, ON, L5L 1J2


Manufactured by: Haupt Pharma Amareg GmbH, Donaustauer Str. 378, Regensburg, Bavaria D-93055, Germany

**Instructions for Use**  
**Natesto (Na-tes-to) CIII**  
**(testosterone)**  
**nasal gel**

**Supplies needed to give your Natesto dose:**

- 1 Natesto dispenser
- 2 clean, dry tissues
- a clean and flat surface, like a table
- a mirror

**Parts of your Natesto dispenser (See Figure A)**



(Figure A)

**Information about your Natesto:**

- **Do not** blow your nose or sniff for 1 hour after using Natesto.
- **Before you use Natesto for the first time, you will need to prime your Natesto pump.**

### How to prime your Natesto pump:

- Hold your Natesto dispenser over a sink, turn it upside down (invert) and slowly press the pump and release 10 times (See Figure B).



(Figure B)

- All the Natesto that comes out when you prime your pump should be rinsed down the sink with warm water.
- If there is any Natesto on the tip of your Natesto actuator after priming, wipe the tip with a clean, dry tissue.
- If any Natesto gel gets on your hands, it is recommended to wash your hands with warm water and soap.

### Giving your Natesto dose:

**Step 1:** Blow your nose.


**Step 2:** Remove the Natesto cap.

**Step 3:** While looking in the mirror, put your right first (index) finger on the pump of your Natesto actuator and slowly slide the tip of the actuator up into your left nostril until your finger on the pump touches the bottom of your nose (See Figure C).



(Figure C)

**Step 4:** Gently tilt the Natesto actuator so that the hole in the tip touches the lateral (side) wall of your nostril. This will make sure that Natesto is given in the correct place (See Figure D).



(Figure D)


**Step 5:** With the actuator in place, slowly push the pump down until it stops and remove the actuator from your nose.

**Step 6:** While looking in the mirror, put your left first (index) finger on the pump of your Natesto actuator and slowly slide the tip of the actuator up into your right nostril until your finger on the pump touches the bottom of your nose (See Figure E).



(Figure E)

**Step 7:** Gently tilt the tip of the Natesto actuator so that the hole in the tip touches the lateral (side) wall of your nostril. This will make sure that Natesto is given in the correct place (See Figure F).




(Figure F)

**Step 8:** Slowly push down on the pump until it stops and remove the actuator from your nose.

**Step 9:** Wipe the tip of the actuator with a clean, dry tissue.


**Step 10:** Replace the cap.

**Step 11:** Press your nostrils together just below the middle of your nose (bridge) and lightly rub them together (See Figure G).



(Figure G)

- Replace your Natesto dispenser when the top of the piston inside the dispenser reaches the arrow at the top of the inside label (See Figure H).



(Figure H)

**How to throw away your empty Natesto dispenser:**

- Safely throw away your empty Natesto dispenser in your household trash.
- Keep Natesto and all medicines out of the reach of children and pets.

This Patient Information and Instructions for Use have been approved by the U.S. Food and Drug Administration.

Marketed by: Acerus Pharmaceuticals Corporation

Manufactured by: Haupt Pharma Amareg GmbH, Donaustauer Str. 378, Regensburg, Bavaria D-93055, Germany

Approved: 10/2016