

TESTOSTER OWN[®]



Actor portrayal.

NATESTO[®] may safely raise testosterone (T) levels while lowering your risk of losing **YOUR OWN T PRODUCTION**

natesto[®] 
(testosterone) nasal gel 

Natesto is a prescription medicine that contains testosterone and is used to treat adult males who have low or no testosterone due to certain medical conditions.

IMPORTANT SAFETY INFORMATION

Who should not use Natesto?

Do not use Natesto if you:

- Are a man who has breast cancer
- Have or might have prostate cancer
- Are pregnant or may become pregnant or are breastfeeding. Natesto may harm your unborn or breastfeeding baby

Talk to your doctor before using Natesto if you have any of the above conditions.

Please see additional Important Safety Information on pages 12-14 and Full Prescribing Information and Patient Information.

Raise your T levels while lowering your risk of losing YOUR OWN T PRODUCTION

For men with T deficiency, also known as hypogonadism, T therapy, such as Natesto, is often recommended.

Like other T therapies, Natesto may help raise T levels and provide symptom relief. However, phase 3 and phase 4 studies suggest that Natesto may also lower the risk of losing your own T production.*†

How might this work? Well, Natesto raises T levels each time you take it (3 times per day). Then, between doses, your T levels return to their pre-dose levels. This gives the body a break from T therapy, while still providing symptom relief. It is thought that this break may help limit the impact on a man's reproductive hormones and natural T production.*

*Natesto is used to treat adult males who have low or no testosterone due to certain medical conditions, and is not indicated for any other health condition.

†T production was assessed by measuring T levels and hormones in a phase 3 study of 306 men and a phase 4 study of 60 men. Sperm count was also measured in the phase 4 study.

IMPORTANT SAFETY INFORMATION (cont)

Natesto may cause serious side effects, including:

- **Changes in red blood cells**
- **Blood clots in your legs or lungs.** Signs and symptoms of a blood clot in your legs can include leg pain, swelling, or redness. Signs and symptoms of a blood clot in your lungs can include difficulty breathing or chest pain
- **Possible increased risk of heart attack or stroke**
- **In large doses, Natesto may lower your sperm count**

Natesto is T therapy that's YOUR OWN IN MORE WAYS THAN ONE



- Raises T levels and provides symptom relief**

See page 6



- Not associated with some of the safety concerns sometimes linked to T therapy

See pages 4 and 5



- May help you maintain reproductive hormones and keep making your own T**†

See page 2



- Doesn't easily transfer to loved ones and others

See page 4



- Is self-administered—no appointments are needed to receive treatment

See page 8

**Shown in a phase 3 study of 306 men. Measuring T levels was the main goal of the study; measuring symptom relief was a secondary goal.

Please see additional **Important Safety Information** on pages 12-14 and [Full Prescribing Information](#) and [Patient Information](#).

Understanding Natesto SAFETY INFORMATION

Natesto has a low rate of overall side effects. Plus, it has a low risk of certain side effects that are sometimes linked to T therapy.



LOW TRANSFERENCE RISK*

- Transferring T to loved ones may be a concern for men who use topical T gels
- With Natesto, no-touch intranasal application reduces the risk of T transference to female partners and children

*Natesto is used to treat adult males who have low or no testosterone due to certain medical conditions, and is not indicated for any other health condition.

Please see the Important Safety Information and Full Prescribing Information for a complete list of side effects associated with Natesto.

IMPORTANT SAFETY INFORMATION (cont)

Natesto may cause serious side effects, including:

- **Changes in red blood cells**
- **Blood clots in your legs or lungs.** Signs and symptoms of a blood clot in your legs can include leg pain, swelling, or redness. Signs and symptoms of a blood clot in your lungs can include difficulty breathing or chest pain
- **Possible increased risk of heart attack or stroke**



LOW RISK OF INCREASING RED BLOOD CELLS*†

- An increase in red blood cells, which may lead to blood clots, is a common side effect of some T therapies
- In a phase 3 study of 306 men, no patients stopped treatment with Natesto due to an increase in red blood cells



NO WARNING FOR BLOOD PRESSURE INCREASES*

- Some T therapies have a warning for increases in blood pressure
- In a phase 3 study of 306 men, no patients stopped treatment with Natesto due to an increase in blood pressure

*Your healthcare provider will check your red blood cell count before and during treatment with Natesto.

IMPORTANT SAFETY INFORMATION (cont)

- **In large doses, Natesto may lower your sperm count**

Please see additional Important Safety Information on pages 12-14 and [Full Prescribing Information](#) and [Patient Information](#).

Natesto raised T levels to **NORMAL RANGE**

In a phase 3, 90-day study of 306 men with T deficiency, the main goal was to evaluate T levels.



Improvements in **SEXUAL FUNCTION AND MOOD SYMPTOMS**

A secondary goal of the phase 3 study was to examine how Natesto affects T deficiency symptoms.

As early as 30 days after starting treatment with Natesto, the study showed some improvements in*:

- **Erectile function**
- **Sexual desire**
- **Intercourse satisfaction**
- **Overall satisfaction**
- **Orgasmic function**
- **Positive and negative feelings**

*Natesto is used to treat adult males who have low or no testosterone due to certain medical conditions, and is not indicated for any other health condition.



IMPORTANT SAFETY INFORMATION (cont)

Before using Natesto, tell your doctor about all of your medical conditions, including if you:

- Have or have had nose (nasal) or sinus problems or nasal or sinus surgery

Nasal application with Natesto was **PREFERRED BY A MAJORITY OF MEN**

In the phase 3 study, 66 men who used Natesto for 90 days took a survey.



IMPORTANT SAFETY INFORMATION (cont)

- Have had a broken nose (fracture) within the past 6 months
- Have or have had a fracture of your nose that caused the inside of your nose to be crooked (deviated anterior nasal septum)
- Have or have had problems with swelling of the lining of your nose (mucosal inflammatory disorder)

Please see additional Important Safety Information on pages 12-14 and [Full Prescribing Information and Patient Information](#).

Innovative, drip-free NASAL APPLICATION

Natesto is a no-drip gel. Each application takes about 10 seconds, applying a thin layer of T that absorbs into the lining of your nose.



—
One pump (5.5 mg) in each nostril 3 times daily, for a total daily dose of 33 mg



—
Each dose should be taken 6 to 8 hours apart: once in the morning, once in the afternoon, once in the evening



—
Try to use Natesto at the same times each day

IMPORTANT SAFETY INFORMATION (cont)

Before using Natesto, tell your doctor about all of your medical conditions, including if you (cont):

- Have breast cancer
- Have or might have prostate cancer
- Have urinary problems due to an enlarged prostate
- Have heart problems
- Have liver or kidney problems
- Have problems breathing while you sleep (sleep apnea)

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, or herbal supplements.

LEARN HOW to use Natesto

Most men are confident applying Natesto within 2 days. To start, prime the pump, then follow 4 simple steps.



PRIME

When starting a new dispenser, be sure to prime the pump before the first use:

- Hold the dispenser upside down and push down 10 times over the sink, then dry the tip
- Safely wash away any dispensed gel. If any gel gets on your hands, wash them with warm water and soap

No priming is needed for subsequent uses of that pump.

—
TIP: It's time for a new dispenser when the top of the piston inside reaches the arrow at the top of the inside label. Reveal the inside label by unwrapping the outer flap from around the container.

GO TO
STEP 1 →

Please see additional Important Safety Information on pages 12-14 and [Full Prescribing Information and Patient Information](#).

Once your dispenser is primed, follow these steps for each dose of Natesto.



1. PREPARE

- Blow your nose
- Remove the dispenser cap



2. POSITION

- While looking in the mirror, place your right index finger on the pump and slowly slide the tip inside your left nostril
- Gently tilt the dispenser so that the hole in the tip touches the soft side of your nostril. This helps ensure Natesto is applied in the correct place

TIP: Your index finger should be touching both the pump and the base of your nose.

IMPORTANT SAFETY INFORMATION (cont)

How should I use Natesto?

- Use Natesto exactly as your doctor tells you to. Your doctor will tell you how much Natesto to use and when to use it
- Natesto is for use in your nose (intranasally) only
- Natesto can be used 30 minutes after using a type of nasal spray called sympathomimetic decongestants such as oxymetazoline. Natesto should not be used with other nasal sprays. It is not known if Natesto is safe and effective when used with other nasal sprays



3. PUMP

- Slowly push down on the pump until it stops. When removing, keep the tip touching the soft side of your nostril. This helps ensure the full dose is applied
- Repeat in your right nostril using your left index finger

TIP: Be sure to completely depress the pump 1 time in each nostril to receive your total dose each morning, afternoon, and evening. Natesto should not be administered in other parts of the body.



4. PRESS

- Wipe the dispenser tip with a clean, dry tissue and replace the cap
- Press your nostrils together just below the bridge of your nose and lightly massage for a few seconds

TIP: Do not blow your nose or sniff for 1 hour after using Natesto.

Please see additional Important Safety Information on pages 12-14 and [Full Prescribing Information and Patient Information](#).

IMPORTANT SAFETY INFORMATION

INDICATION

What is Natesto Nasal Gel?

Natesto is a prescription medicine that contains testosterone and is used to treat adult males who have low or no testosterone due to certain medical conditions. Your doctor will test the testosterone level in your blood before you start and while you are using Natesto.

Natesto should not be used by men with breast cancer, or known or suspected prostate cancer.

It is not known if Natesto is safe or effective to treat men who have low testosterone due to aging.

Safety and effectiveness of Natesto have not been established in children younger than 18 years old, and Natesto should not be used by children. Natesto use by children may affect bone growth.

Natesto is a controlled substance (CIII) because it contains testosterone that can be a target for people who abuse prescription medicines. Keep your Natesto in a safe place to protect it. Never give your Natesto to anyone else, even if they have the same symptoms you have. Selling or giving away this medicine may harm others and is against the law.

Natesto is not meant for use in women.

IMPORTANT SAFETY INFORMATION

Who should not use Natesto?

Do not use Natesto if you:

- Are a man who has breast cancer
- Have or might have prostate cancer
- Are pregnant or may become pregnant or are breastfeeding. Natesto may harm your unborn or breastfeeding baby

Talk to your doctor before using Natesto if you have any of the above conditions.

What should I tell my doctor before using Natesto?

Before using Natesto, tell your doctor about all of your medical conditions, including if you:

- Have or have had nose (nasal) or sinus problems or nasal or sinus surgery
- Have had a broken nose (fracture) within the past 6 months
- Have or have had a fracture of your nose that caused the inside of your nose to be crooked (deviated anterior nasal septum)
- Have or have had problems with swelling of the lining of your nose (mucosal inflammatory disorder)
- Have breast cancer

- Have or might have prostate cancer
 - Have urinary problems due to an enlarged prostate
 - Have heart problems
 - Have liver or kidney problems
 - Have problems breathing while you sleep (sleep apnea)
- Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, or herbal supplements.

Using Natesto with certain other medicines can affect each other. Especially tell your doctor if you take:

- Insulin
- Medicines that decrease blood clotting
- Corticosteroids

How should I use Natesto?

- Use Natesto exactly as your doctor tells you to. Your doctor will tell you how much Natesto to use and when to use it
- Natesto is for use in your nose (intranasally) only
- Natesto can be used 30 minutes after using a type of nasal spray called sympathomimetic decongestants such as oxymetazoline. Natesto should not be used with other nasal sprays. It is not known if Natesto is safe and effective when used with other nasal sprays

What are the possible side effects of Natesto?

Natesto may cause serious side effects, including:

- **Nose (nasal) problems.** Signs and symptoms of nasal problems may include runny nose, congestion, sneezing, nosebleed, nasal discomfort, nasal scabbing, or nasal dryness
- **If you already have enlargement of your prostate gland, your signs and symptoms can get worse while using Natesto.** This can include: increased urination at night, trouble starting your urine stream, having to pass urine many times during the day, having an urge that you have to go to the bathroom right away, having a urine accident, and being unable to pass urine or weak urine flow
- **Possible increased risk of prostate cancer.** Your doctor should check you for prostate cancer or any other prostate problems before you start and while you use Natesto

Please see additional Important Safety Information on page 14 and [Full Prescribing Information](#) and [Patient Information](#).

IMPORTANT SAFETY INFORMATION (cont)

Natesto may cause serious side effects, including (cont):

- **Changes in red blood cells**
- **Blood clots in your legs or lungs.** Signs and symptoms of a blood clot in your legs can include leg pain, swelling, or redness. Signs and symptoms of a blood clot in your lungs can include difficulty breathing or chest pain
- **Possible increased risk of heart attack or stroke**
- **In large doses, Natesto may lower your sperm count**
- **Possible increased risk of liver problems.** Signs and symptoms of liver problems may include: nausea or vomiting, yellowing of your skin or whites of your eyes, dark urine, or pain on the right side of your stomach area (abdominal pain)
- **Swelling of your ankles, feet, or body, with or without heart failure.** This may cause serious problems for people who have heart, kidney, or liver disease
- **Enlarged or painful breasts**
- **Problems breathing while you sleep (sleep apnea)**

Call your doctor right away if you have any of the serious side effects listed above.

The most common side effects of Natesto include:

- Increased prostate-specific antigen (a test used to screen for prostate cancer)
- Headache
- Runny nose
- Nosebleeds
- Nose pain
- Sore throat
- Cough
- Upper respiratory infection
- Sinus infection
- Nasal scabs

Other side effects include more erections than normal for you or erections that last for a long time.

These are not all the possible side effects of Natesto. Call your doctor for medical advice about side effects.

Store Natesto between 68°F to 77°F (20°C to 25°C).

Keep Natesto and all medicines out of the reach of children.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

For more information, visit Natesto.com, or call 1-833-698-3786.

Get savings and support with Natesto

Natesto Savings Program



Eligible, commercially insured patients may **pay as little as \$0 per month** for Natesto, for a total savings of up to \$3500 per year.*

Natesto Cash Option Program



Actor portrayal.

Cash patients and insured, not covered patients may fill each valid Natesto prescription or refill at the **cash price of \$140.***

You can also
**CONNECT BY
SCANNING
THIS CODE**



*See full eligibility requirements and restrictions on back.

Please see additional **Important Safety Information** on pages 12-14 and **Full Prescribing Information and Patient Information.**

Natesto Savings Program Terms and Conditions

The Natesto Savings Program is valid ONLY for patients with private (commercial) insurance. Patients who are enrolled in any federal healthcare program, including, without limitation, Medicare Part D, Medicare Advantage, Fee-for-Service Medicaid, Managed Medicaid, Veterans Affairs, Champus, TRICARE, and Indian Health Service, are not eligible to participate in the Natesto Savings Program. Eligible, commercially insured patients may pay as little as \$0 co-pay on each valid Natesto prescription or refill, up to an annual maximum of \$3500. Once a patient exceeds the \$3500 annual benefit in the plan or payer's benefit year, the patient is responsible for all remaining co-pay or cost-sharing obligations throughout the conclusion of the current benefit year. On each valid Natesto prescription or refill, the Natesto Savings Card benefit may be applied to dispensed quantities of no less than 20 days (2 dispensers) to no more than 90 days (9 dispensers) per prescription dispensed. **The Natesto Savings Program is not health insurance.**

Some restrictions may apply due to state regulations. Patients are responsible for reporting receipt of this coupon benefit to any insurer, health plan, or other third party who pays for or reimburses any part of the prescription filled using the coupon, as may be required by payer or plan rules, or in accordance with any state or federal law. Pharmacists, patients, and prescribers agree not to seek reimbursement for all or any part of the benefit received by the patient through this offer from any third-party payer or from any health savings, flexible spending, or other healthcare reimbursement account.

The Natesto Savings Program offer is valid only for US male residents 18 years of age or over. Offer good only in the USA and void where prohibited by law, taxed, or restricted. Acerus Pharmaceuticals Corporation reserves the right to rescind, revoke, or amend this offer without notice. Card is limited to one per person, is not transferable, and cannot be reproduced. For further information or assistance with the Natesto Savings Program, please call 1-855-390-0162.

Natesto Cash Option Program Terms and Conditions

Cash patients and insured, not covered patients may fill each valid Natesto prescription or refill at the cash price of \$140 for quantities of no less than 20 days (2 dispensers), and not to exceed 30 days (3 dispensers), per prescription dispensed for up to 12 refills. The Natesto Cash Option Program does not process insurance claims. Additionally, commercially insured patients who exceed the \$3500 out-of-pocket maximum for the Natesto Savings Card benefit program and have a co-pay or co-insurance requirement that exceeds \$140 per 30-day prescription in a given 12-month benefit year, according to payer or plan design, may opt into the Natesto Cash Option Program until the new benefit year begins. Patients may not apply the Natesto Cash Option Program payments to any deductible, co-pay, or to any other out-of-pocket spending calculations for any insurer or third-party payer, including any government healthcare program, such as Medicare Part D true out-of-pocket (TrOOP) costs. **The Natesto Cash Option Program is not health insurance.**

The Natesto Cash Option Program is valid only for US male residents 18 years of age or over. The Natesto Cash Option Program may be filled through a retail pharmacy. Patients are responsible for reporting use of the Natesto Cash Option Program to any private insurer, health plan, or third-party payer. Acerus Pharmaceuticals Corporation reserves the right to rescind, revoke, or amend this offer without notice, at any time, at its sole discretion. For further information or assistance with the Natesto Cash Option Program, please call 1-855-390-0162.

**Please see Important Safety Information on pages 12-14
and [Full Prescribing Information](#) and [Patient Information](#).**

